

ORIGINAL RESEARCH

Prevalence of presbyopia in a rural African community

JN Uche, IR Ezegwui, E Uche, EN Onwasigwe, RE Umeh, CN Onwasigwe

Department of Ophthalmology, University of Nigeria Teaching Hospital, Enugu, Nigeria

Submitted: 28 June 2013; **Revised:** 20 November 2013; **Accepted:** 11 December 2013; **Published:** 6 August 2014

Uche JN, Ezegwui IR, Uche E, Onwasigwe EN, Umeh RE, Onwasigwe CN

Prevalence of presbyopia in a rural African community

Rural and Remote Health 14: 2731. (Online) 2014

Available: <http://www.rrh.org.au>

ABSTRACT

Introduction: When presbyopia (loss of accommodation of the crystalline lens with increasing age) sets in, doing near work becomes associated with headache and eye strain. Reading and writing become a challenge. Literacy levels may be low in rural communities; nevertheless some work other than reading, like sewing, sorting stone from grain and operating mobile phones, is done with dissatisfaction. This study aims to determine the prevalence of presbyopia, the unmet presbyopia need and the presbyopia correction coverage in a rural African community.

Methods: A population-based cross-sectional study was carried out in a rural population aged 35 years and greater, selected by cluster random sampling. Information was sought on biodata of the participants and they were subsequently examined. Distance visual acuity for each participant was determined. Anterior and posterior segments of the eyes were examined. Objective refraction with subjective refinement was done on all subjects with distant visual acuity less than 6/6. Near visual acuity was assessed at 40 cm with distant correction in place if required. Presbyopia was defined as inability to read N8 at 40 cm or requiring an addition of at least +1.00DS to improve near vision to at least N8. Questionnaires were administered to those identified as presbyopic on source of procurement of spectacles (if they had one) and on reasons for non-procurement of presbyopic spectacles. They were also asked to rate their difficulty with various listed near work. Data entry and analysis were done using Statistical Package for the Social Sciences v16.0 and Program for Epidemiologist v4.01 software.

Results: A total of 585 subjects (participation rate 81.1%) aged 35 years and greater were interviewed and examined. The prevalence of presbyopia was 63.4% (95% confidence interval (CI) 62.6–64.2%). There was increasing prevalence with increasing age. The met presbyopia need was 17.6%, unmet need was 45.8% and presbyopic correction coverage was 27.8%. The commonest reasons for not procuring presbyopic correction were 'not a priority' (21.5%) and 'cost' (21.2%).



Conclusions: The prevalence of presbyopia in this rural African community is high. Many who need presbyopic correction do not have corrective spectacles.

Key words: Nigeria, presbyopia, presbyopic correction coverage, prevalence, rural community.

Introduction

The World Health Organization estimates that 153 million people are blind or visually impaired from uncorrected refractive error¹. This does not include the impairment of near vision from uncorrected presbyopia. Holden et al², using multiple population-based surveys, estimated that 1.04 billion people globally have presbyopia. Population-based surveys from rural Tanzania³, South India⁴, Brazil⁵ and Iran⁶ recorded presbyopia prevalences of 62%, 55.3%, 54.7% and 58.2% respectively. The minimum age of the study subjects was not uniform in these studies.

Presbyopia is the loss of accommodation of the crystalline lens with increasing age. From early childhood the amplitude of accommodation is about 15 dioptres (D) and this continues to decrease progressively to about 1D by the age of 60 years. With failing accommodation, doing near work becomes associated with headache and eye strain. Reading and writing become a challenge. Even in rural communities, where literacy levels may be low, dissatisfaction occurs when presbyopia sets in⁷, because the condition affects near work like sewing, sorting rice and winnowing grain. Moreover, mobile phones are increasingly being used in rural communities, unmasking the need to address the presbyopia challenge there.

Previous studies from low- and middle-income countries⁷ suggest that more than half of adults aged 30 years and greater have presbyopia, with women being more affected both in prevalence and in severity. Most of these individuals do not have corrective spectacles. Education and literacy have been identified as important tools in achieving Millennium Development Goals⁸. With the emphasis on adult education and literacy, it becomes necessary to identify the level of

presbyopia in the population with a view to developing refraction services to correct them. This study, therefore, aims to determine the prevalence of presbyopia, the unmet presbyopia need and the presbyopia correction coverage in a rural African community.

Methods

Study area

Enugu East Local Government Area is one of the 17 local government areas that make up Enugu state, south-eastern Nigeria. It has a population of 259 924. It is divided into four zones: Abakpa, Nike, Trans-Ekulu and Emene. Nike has a total population of 51 411, and it is mostly rural. The residents are predominantly subsistence farmers. Six villages make up Nike. A population based cross-sectional study was carried out between June and July 2010 in this community, located about 20 km away from Enugu, the capital city.

Study population

The study population is adults 35 years and older residing in Nike. The minimum sample size was calculated using the formula for sample size for a definite population. The sample size was calculated to be 720, considering 53% prevalence from a local study⁹ as the proportion of the population with presbyopia and assuming 0.05 as the sampling error, 95%CI with a design effect of 1.5, allowing 20% for non-response.

The cluster random sampling method with probability proportional to size was used. Eight study clusters were selected and 90 individuals aged 35 years and greater were enumerated from each cluster. A cluster is made up of several



households demarcated by major footpaths. An enumerator from the community carried out the enumeration of the households and subjects after adequate training.

Methods

Pre-survey visits were made to the leaders of the communities to familiarise them and the community with the survey. They assigned a guide and an enumerator who worked with the research team.

Written informed consent was obtained from each participant prior to being interviewed and examined. The interviews and examinations took place in health centres in the communities that had them; otherwise, a classroom in the community's school was used. Two research assistants who were trained for the survey interviewed the participants using a questionnaire designed for the study. Information was sought on demographic data and subjects were then examined. Those identified as having presbyopia were asked further questions relating to where they procured their glasses from (if they had them) and barriers to obtaining presbyopic correction. They were asked to rate the difficulty they had with doing various listed near work on a scale with choices 'none', 'little', 'moderate' or 'great'.

One ophthalmologist (NJU) examined and refracted all the participants. Visual acuity (VA) with and without pinhole was tested using Snellen's chart at 6 m. Anterior segment examination was done with torch light and head loupe. The posterior segment was examined with a direct ophthalmoscope.

Refraction was carried out by streak retinoscopy on all subjects with VAs less than 6/6. The best corrected visual acuity (BCVA) was recorded. Near visual acuity was measured binocularly using the N notation illiterate E chart at a distance of 40 cm from the eyes. An inextensible string was attached to the near vision chart and the other end placed against the subject's forehead and held firmly to ensure a distance of 40 cm from the eye. Spherical plus lenses were added using increments of 0.25D until the patient could read

N8. Subjects with VAs of 6/6 were assumed emmetropic and tested for near vision as described above.

Subjects with reduced vision that could not be improved by refraction and those with additional ocular pathology were referred to the tertiary eye care centres in the state. Simple ocular conditions like bacterial and allergic conjunctivitis were treated on site at no cost to the subjects. Reading glasses were given free of charge to those who needed them.

Data analysis

All data were cleaned, double-entered into the computer and analyzed using Statistical Package for the Social Sciences v16.0 (SPSS; <http://www.spss.com>) and Program for Epidemiologist v4.01 software (<http://www.brixtonhealth.com/pepi4windows.html>). Frequency and contingency tables were used to present the distribution of data. The χ^2 test as a statistical test of significance was done for the discrete variables. Multivariate analysis in the form of logistic regression was used to predict the factors that are associated with presbyopia. All statistical calculations were done at the significance of $p < 0.05$.

Inclusion and exclusion criteria

All individuals resident in the community aged 35 years and above were eligible to be enrolled.

Exclusion criteria were:

1. Age less than 35 years.
2. Non-residence in the community. A regular resident is defined as somebody who has lived in the study area continuously for the previous 6 months.
3. Visual acuity worse than 6/60 in the better eye, not improving with pinhole.

Definitions

Presbyopic: could not read the N8 optotype at 40 cm with the distance correction in place (if required). (N8 was chosen because it is the size of the local newspaper print.)



Functional presbyopia: need for a significant optical correction ($\geq +1.00D$) added to the presenting distance refractive correction to achieve a near visual acuity criterion of N8 print².

Objective presbyopia: need for a significant optical correction ($\geq +1.00D$) added to the best distance optical correction to improve near vision to a near visual acuity criterion of N8².

Met presbyopia need (MPN): measure of dispersion of spectacles to correct those with presbyopia to see N8 or better binocularly¹⁰.

$$MPN = 100 \times \text{number who see N8 or better with own near vision spectacles} \div \text{total sample population}$$

Unmet presbyopia need (UPN): number of subjects unable to see N8 binocularly¹⁰.

$$UPN = 100 \times \text{number unable to see N8 binocularly} \div \text{total sample population}$$

Presbyopic correction coverage (PCC): $MPN \div (MPN + UPN) \times 100^{10}$.

Ethics approval

Ethics clearance was obtained from the Health Research Ethics Committee of University of Nigeria Teaching Hospital Enugu (approval numbers are not issued).

Results

Of the 720 subjects enumerated, 585 took part in the study (81.1% participation rate). Eighty seven (12%) were not available on the examination days while 48 (7%) were not eligible based on exclusion criteria. No subject refused to participate.

The mean age ($\pm$ standard deviation) of the subjects was 49 (± 11.1) years. The age range was 35–99 years. Demographic characteristics are shown in Table 1. The majority were manual workers (277, 47.4%) and had little or no formal education (375, 64.1%). The prevalence of objective presbyopia was 63.4% (95%CI 62.6–64.2; Table 2). Prevalence of presbyopia increased with increasing age. The mean severity of presbyopia in dioptres was higher for females than males aged 35–54 years but this reverses for the age group 55 years and older. Logistic regression showed that occurrence of presbyopia was more likely as one gets older. Age was the most predictive factor and the associate was statistically significant ($p < 0.01$; Table 3).

Of the 371 presbyopes, 103 had reading spectacles while 268 had none. This gives an MPN of 17.6%, UPN of 45.8% and presbyopia coverage of 27.8% (Table 4). There was a statistically significant difference in the gender, educational level and occupation of the subjects with MPN and those with UPN ($p < 0.01$, 0.001 and 0.001 respectively). This is reflected in the presbyopia correction coverage. Thus the presbyopia coverage was higher for males than females, for those with tertiary or secondary education than primary or no formal education, and for skilled workers more than manual workers.

Among the presbyopic subjects who had spectacles, 47.6% obtained their spectacles from an eye clinic, 33% from an optical shop, while the rest (19.4%) obtained theirs from the market (19) or outreach (1).

The major reasons for not purchasing reading glasses were 'not a priority' (131, 21.5%), 'cost' (129, 21.2%) and 'not aware of the problem' (108, 17.8%). Others were 'normal aging process' (34, 5.6%), 'eye care services too far' (21, 3.5%), 'don't know where to go' (20, 3.3%), 'lost or broke glasses' (7, 1.2%), 'no companion' (5, 0.8%) and miscellaneous reasons (153, 25.2%). Table 5 shows the proportion of the subjects who had difficulty with various near vision tasks. Most participants experienced difficulty with reading and threading a needle.



Table 1: Distribution of sample according to baseline characteristics

Characteristic	Male (n=191) No. (%)	Female (n=394) No. (%)	χ^2 test	p value
Age group (years)				
35–44	38 (19.9)	173 (43.9)	36.842	<0.001***
45–54	78 (40.8)	136 (34.5)		
55–64	42 (22.0)	50 (12.7)		
≥65	33 (17.3)	35 (8.9)		
Education level				
None	34 (17.8)	146 (37.1)	22.833	<0.001***
Primary	74 (38.7)	121 (30.7)		
Secondary	36 (18.8)	60 (15.2)		
Tertiary	47 (24.6)	67 (17.0)		
Occupation				
Unemployed	19 (9.9)	27 (6.9)	9.709	0.021*
Manual	73 (38.2)	204 (51.8)		
Skilled	33 (17.3)	55 (14.0)		
Other	66 (34.6)	108 (27.4)		

* $p < 0.05$, *** $p < 0.001$

Table 2: Prevalence of presbyopia by gender and age group

Age group (years)	Male (n=191) No.	Presbyopic male No. (%)	Female (n=394) No. (%)	Presbyopic female No. (%)	χ^2 test	p value
35–54	38	12 (31.6)	173	103 (59.5)	39.362	<0.001***
45–54	78	56 (71.8)	136	88 (64.7)		
55–64	42	29 (69.0)	50	33 (66.0)		
≥65	33	24 (72.7)	35	26 (74.3)		

*** $p < 0.001$

Table 3: Logistic regression to determine factors associated with presbyopia

Factor	Odds ratio	p value
Age	1.033	0.001**
Gender	1.097	0.650
Education	0.863	0.105
Occupation	1.003	0.967

** $p < 0.01$

Discussion

The prevalence of presbyopia in this study was 63.4%. Nwosu¹¹, who also worked in south-eastern Nigeria,

examined young adults 18–49 years and found a prevalence of 33%. Marmamula et al¹² found a prevalence of 63.7%, which is similar to the finding in this study.



Table 4: Presbyopic correction coverage, presbyopic met need and unmet need by subject characteristic

Characteristic	Sample (n=585)	Met presbyopia need No. (%)	Unmet presbyopia need No. (%)	Presbyopia correction coverage (%)	χ^2 -test	p value
Age group (years)						
35–44	211	31 (14.7)	84 (39.8)	27.0	2.515	0.473
45–54	214	46 (21.5)	98 (45.8)	31.9		
55–64	92	15 (16.3)	47 (51.1)	24.2		
≥65	68	11 (16.2)	39 (57.4)	22.0		
Gender						
Male	191	47 (24.6)	74 (38.7)	38.8	10.992	0.001**
Female	394	56 (14.2)	194 (49.2)	22.4		
Education						
None	180	8 (4.4)	112 (62.2)	6.7	73.391	<0.001***
Primary	195	34 (17.4)	103 (52.8)	24.8		
Secondary	96	23 (24.0)	33 (34.4)	41.1		
Tertiary	114	38 (26.4)	20 (17.5)	65.5		
Occupation						
Unemployed	46	9 (19.6)	24 (52.2)	25.0	38.604	<0.001***
Manual labour	277	31 (11.2)	156 (56.3)	16.6		
Skilled labour	88	29 (33.0)	19 (21.6)	60.4		
Other	174	34 (19.5)	69 (39.7)	33.0		

** $p < 0.01$, *** $p < 0.001$

Table 5: Proportion of sample reporting difficulty with near visual tasks without previous spectacles

Visual task	Number	Mean score	Standard deviation
Reading	585	2.15	1.25
Writing	585	1.57	1.42
Cooking food	585	0.12	0.48
Winnowing	585	0.21	0.66
Sorting out rice	584	0.41	0.93
Threading a needle	585	2.18	1.20
Cutting fingernails	584	0.47	0.98
Dressing children	583	0.05	0.32
Weeding	582	0.07	0.36
Harvesting	584	0.08	0.42
Recognizing faces close by	583	0.11	0.48
Recognizing small objects	580	0.24	0.71

Great difficulty = 3; moderate difficulty = 2; little difficulty = 1; no difficulty = 0

Mean score ≥ 2 = problem; mean score < 2 = no problem

The varying differences in prevalence of presbyopia in studies from low- and middle-income countries may arise from different definitions of presbyopia, different minimum age of study subjects and different examination conditions (outdoors or indoors). Some studies¹⁰ examined for functional presbyopia while others examined for objective presbyopia⁴.

This study examined for objective presbyopia. When objective presbyopia is used, people with low or moderate myopia are more likely to be identified as needing presbyopic correction so the estimated prevalence will be higher. The mean severity of presbyopia in the present series was higher in females aged less than 55 years but higher in males above



this age. The reason for this could not be ascertained. Some authors^{4,6,7} documented an increase in severity and prevalence of presbyopia in females across all age groups.

The presbyopic correction coverage, although low, is higher in males, skilled workers and those with some education. This is similar to the findings in Timor-Leste¹³. The major reasons for not purchasing spectacles were 'not a priority' and 'cost'. Some authors¹⁴ found 'lack of felt need' and 'awareness' to be the important barriers to having presbyopic correction in rural Andhra Pradesh, India. 'Lack of felt need' and 'not a priority' are somewhat similar. In rural China¹⁵ 'concerns about poor quality of available correction' (33%) and 'lack of awareness' (29%) were the commonest reasons. In a rural setting where poverty abounds and feeding is a struggle, procuring presbyopic corrections is not rated a priority. The subjects do not realize that correcting their near vision impairment can increase their efficiency at doing near work (for example sewing and hairdressing). Their quality of life can also be improved by correction of their near vision impairment. Mobile phones are increasingly being used in rural communities in Nigeria and require good near vision to operate well. Articles of worship like hymn books are also widely used in these communities. There is a need to educate the people in these rural communities about the importance of correction of near visual impairment. Thereafter refraction services can be integrated into primary eye care in their communities. Affordable spectacles can then be made available to them.

A limitation of this study was that some of the participants were not sure of their date of birth. Historic events were used to estimate their age.

Conclusions

The prevalence of presbyopia in this study was high. Many individuals with near vision impairment did not have any presbyopic correction. There is a need to address the problem of presbyopia in this rural community through health education, to make residents consider correction of presbyopia a priority.

References

1. World Health Organization. *Vision 2020 global initiative for the elimination of avoidable blindness*. Fact sheet no 1231. Geneva: WHO, 2000.
2. Holden BA, Fricke TR, Ho SM, Wong R, Schlenther G, Cronje S et al. Global vision impairment due to uncorrected presbyopia. *Archives of Ophthalmology* 2008; **126**: 1731-1739.
3. Burke AG, Patel I, Munoz B, Kayongoya A, McHiwa W, Schwarzwaldner AW et al. Population-based study of presbyopia in rural Tanzania. *Ophthalmology* 2006; **113**: 723-727.
4. Nirmalan P, Krishnaiah S, Shamanna BR, Rao GN, Thomas R. A population-based assessment of presbyopia in the state of Andhra Pradesh Eye Disease Study. *Investigative Ophthalmology and Visual Science* 2006; **47**: 2324-2328.
5. Duarte W, Barros A, Dias-da-Costa JS, Cattán JM. Prevalence of near vision deficiency and related factors: a population-based study. *Cadernos de Saude Publica* 2003; **19**: 551-559.
6. Hashemi H, Khabazkhab M, Jafarzadehpour E, Mehravaran S, Emamian MH, Yekta A et al. Population-based study of presbyopia in Sharoud, Iran. *Clinical & Experimental Ophthalmology* 2012; **40**: 863-868.
7. Patel I, West SK. Presbyopia: prevalence, impact and interventions. *Community Eye Health* 2007; **20**: 40-41.
8. Fall H. Vision 2020 and the millennium development goals. *Community Eye Health* 2007; **20**: 62-64.
9. Chiroma M. Prevalence of presbyopia and the impact of uncorrected presbyopia on quality of life in rural Gwagwalda, Abuja. International Centre for Eye Health MSc Dissertation, 2008.
10. Sherwin JC, Keeffe JE, Kuper H, Islam FM, Muller A, Mathenge W. Functional presbyopia in a rural Kenyan population: the unmet presbyopia need. *Clinical & Experimental Ophthalmology* 2008; **36**: 245-251.
11. Nwosu SN. Ocular problems of young adults in rural Nigeria. *International Ophthalmology* 1998; **22**: 259-263.



12. Marmamula S, Keefe JE, Rao GN. Uncorrected refractive errors, presbyopia and spectacle coverage. Results from a rapid assessment of refractive error survey. *Ophthalmic Epidemiology* 2009; **16**: 269-274.

13. Ramke J, du Toit R, Palagyi A, Brian G, Naduviath T. Correction of refractive error and presbyopia in Timor-Leste. *British Journal of Ophthalmology* 2007; **91**: 860-866.

14. Marmamula S, Keefe JE, Raman U, Rao GN. Population-based cross-sectional study of barriers to utilization of refraction services in South India: Rapid Assessment of Refractive Errors (RARE) Study. *British Medical Journal Open* 2011; **1**(1): e000172.

15. Lu Q, Congdon N, He X, Murthy GV, Yang A, He W. Quality of life and near vision impairment due to functional presbyopia among rural Chinese adults. *Investigative Ophthalmology & Visual Science* 2011; **52**: 4118-4123.
