

RESEARCH ARTICLE

Refractive error and its associated factors among pregnant women attending antenatal care unit at the University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia


Mengistie Diress^{1*}, Yigizie Yeshaw^{1,2}, Minychil Bantihun³, Baye Dagne¹, Adugnaw Ambelu¹, Mohammed Abdu Seid⁴, Yonas Akalu¹

1 Department of Human Physiology, School of Medicine, University of Gondar, Gondar, Ethiopia,

2 Department of Epidemiology and Biostatistics, Institute of Public Health, University of Gondar, Gondar, Ethiopia, **3** Department of Optometry, School of Medicine, University of Gondar, Gondar, Ethiopia,

4 Department of Human Physiology, School of Medicine, Debre Tabor University, Debre Tabor, Ethiopia

* mengistiediress@gmail.com



Abstract

OPEN ACCESS

Citation: Diress M, Yeshaw Y, Bantihun M, Dagne B, Ambelu A, Seid MA, et al. (2021) Refractive error and its associated factors among pregnant women attending antenatal care unit at the University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia. PLoS ONE 16(2): e0246174. <https://doi.org/10.1371/journal.pone.0246174>

Editor: Frank T. Spradley, University of Mississippi Medical Center, UNITED STATES

Received: September 24, 2020

Accepted: January 14, 2021

Published: February 12, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0246174>

Copyright: © 2021 Diress et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Background

Refractive error is one of the commonly encountered problems during pregnancy and being the cause of deleterious effects on health. Despite its impacts, there is no evidence on the magnitude and associated factors of refractive error among pregnant women in Ethiopia. This study aimed to determine the prevalence of refractive error and its associated factors among pregnant women attending antenatal care unit at the University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia, 2020.

Methods

An institution-based cross-sectional study was employed. An ocular examination was performed using Retinoscope and Snellen's illiterate "E" chart. The required data were collected using an interviewer-administered questionnaire which comprised socio-demographic, clinical and pregnancy-related variables. EpiData 3.02 and STATA 14 were used for data entry and analysis respectively. Both bivariable and multivariable binary logistic regression analyses were executed to identify factors associated with refractive error. Variables with a p-value ≤ 0.05 in the multivariable logistic regression analysis were declared as significantly associated factors with refractive error.

Results

A total of 401 pregnant women with a median age of 27 (IQR = 24–31) years participated in this study. The overall prevalence of refractive error among the study participants was 35.66% (95% CI: 30.95–40.37). Of the total study participants, ninety-two (22.90%) of them were myopic, forty-five (11.22%) were hyperopic and the rest were antimetropic. Increased

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: ANC, Ante-Natal Care; D, Diopters; GDM, Gestational Diabetes Mellitus; PIH, Pregnancy Induced Hypertension; SE, Spherical Equivalent; UoG, University of Gondar; UoGCSH, University of Gondar Comprehensive Specialized Hospital; WHO, World Health Organization.

maternal age (AOR = 1.31; 95% CI: 1.16–1.48)), increased parity (AOR = 3.17, 95% CI: 1.92–5.25), increased gestational age (AOR = 1.15, 95% CI: 1.08–1.22), and regular use of computers/ watching television (AOR = 6.19, 95% CI: 2.46–15.59) were significantly associated with refractive error.

Conclusion

The prevalence of refractive error among pregnant women was high where myopia was the most common variety. Advanced maternal age, increased gestational age, increased parity and regular use of computer or watching television were significantly associated with refractive error among pregnant women. Therefore, apart from providing other maternal health services, routine screening and evaluation of pregnant women for refractive error during antenatal care visit is recommended to avoid its negative impacts.

Introduction

Refractive error (RE) is the leading cause of visual impairment worldwide [1–3]. Globally, 333 million people are visually impaired. Of these, 153 million cases are attributed to uncorrected refractive error [2,3]. The prevalence of RE is about 46% in Sub-Saharan Africa [3,4]. According to the 2006 National Survey Report of Ethiopia, RE is the second cause of visual impairment (33.40%) following cataract (42.3%) and the third cause of blindness (7.80%) [5].

Studies in the world consistently showed that females have a relatively higher risk of developing RE than males [6,7]. The latest study in Saudi Arabia estimated that the prevalence of RE among women is 25.36% [8]. RE is also a common problem among pregnant women in developing countries [9,10]. A study in India revealed that 65% of pregnant women have myopia [11]. The finding of another study in Nigeria shows that myopia is the most prevalent type of RE (57%) followed by hyperopia (33%) [10]. The prevalence of RE in South India is estimated to be 77.50% [12].

Uncorrected RE among pregnant women is associated with both immediate and long-term consequences. This problem leads to disabilities, unable to travel safely through the environment, loss of educational and job opportunities, and reduced productivity among the women. It can also spoil the quality of life of the women and may give rise to retinal detachment if not treated early [13–17].

Evidences from different regions of the world revealed that RE is associated with maternal age [18,19], residence [20], level of education [21,22], gestational age [10,23,24], mental stress [25], sleep disturbance [26], family history of RE [27], contraceptive use [28], iron deficiency anemia [29], gestational diabetes mellitus (GDM) [30,31], pregnancy induced hypertension (PIH) [32,33], medication history [34], and prolonged use of digital devices [1,35,36].

Though various literatures revealed the existence of numerous pregnancy-associated visual problems, little is known about RE in Africa particularly in Ethiopia. Therefore, this study aimed to determine the prevalence and associated factors of RE among pregnant women attending antenatal care (ANC) unit at the University of Gondar Comprehensive Specialized Hospital (UoGCSH). Conducting this research will have a crucial role in filling the above gaps, and to design and implement strategies for the prevention and management of the problem in the study setting.

Methods and materials

Ethical approval and consent to participate

Ethical clearance with a reference number of 1828/2012 was obtained from the Institutional Review Board (IRB) of the UoG and letter of cooperation was gotten from the UoGCSH. Written informed consent was taken from each study participant. Privacy and confidentiality of information were kept properly and names were not recorded. Those study participants who had severe RE at the time of data collection were referred to the Department of Ophthalmology for further diagnosis and management.

Study setting, design, and population

An institution-based cross-sectional study design was carried out at the UoGCSH, ANC unit from 01 February 2020, to 30 March 2020. The University of Gondar (UoG) is located 727 km far from the capital city, Addis Ababa in the Northwest direction. ANC unit of the UoGCSH is one of the outpatient departments established earlier and currently provides services for about 22,824 pregnant women per year. Pregnant women visiting the ANC service are regularly evaluated and the findings were documented in their charts. All pregnant women who had ANC service at the UoGCSH were included in the study whereas, pregnant women with known diabetes mellitus and hypertension before being pregnant, those with congenital eye problems, refractive problems, glaucoma, cataract, and eye trauma during the study period were excluded from the study.

Sample size determination and sampling procedure

The required sample size for the study was estimated using a single population proportion formula, taking an expected estimate of the prevalence of 50% (no previous study was conducted in Ethiopia), 95% confidence interval, and a 5% margin of error. Accordingly, the sample size for this study became 384.

$$N = \frac{(Z_{\alpha/2})^2 * p(1 - p)}{d^2} = \frac{(1.96)^2 * 0.5(1 - 0.5)}{(0.05)^2} = 384$$

Where; N = sample size, p = expected estimates of prevalence value (50%), d = margin of sampling error tolerated- 5% (0.05), $Z_{\alpha/2}$ = critical value at 95% confidence interval of certainty (1.96).

After adding a 5% non-response rate, the final sample size became 403. Then, a systematic random sampling technique was used to recruit the study participants. The first participant was recruited by lottery method and the remaining participants were recruited by adding 9 (interval size) which was determined by dividing the total number of pregnant women who attended ANC unit during the study period (3804) to the required sample size (403).

Study variables

The dependent variable was refractive error (yes or no). Participants were considered to have a refractive error, if they were either myopic (the spherical equivalent ≤ -0.50 diopters (D)) or hyperopic (the spherical equivalent $\geq +0.50$ D) or both problems in either or both eyes. If the spherical equivalent (SE) was in between -0.50D and +0.50D in either or both eyes, they were considered as normal [37,38].

The independent variables for this study were age, residence, income, occupation, educational level, parity, gestational age, GDM, preeclampsia, and eclampsia, Catha edulis (khat),

cigarette smoking, coffee drink, alcohol intake, anemia, medication history, stress, sleep disturbance, family history of spectacle use, and regular use of computers or watching TV.

Operational definitions

Refractive error. A spherical equivalent (SE) of less than -0.50 and/or greater than $+0.50$ D in either eye or both [37,38].

Myopia. A spherical equivalent (SE) of less than -0.50 D in either eye or both, that can be corrected by concave lens [38,39].

Hyperopia. A spherical equivalent (SE) of greater than $+0.50$ D in either eye or both, that can be corrected by convex lens [38].

Antimetropia. A situation in which one eye is hyperopic, while the other is myopic [40].

Anemia. Hemoglobin level of less than 12.00 g/dl or hematocrit level of less than 36.00% [41].

Sleep disturbance. Sleeping time of ≤ 5 hours/day or sleeping time of ≥ 9 hours/day [42,43].

Substance use. The use of at least one of the substances (alcohol, khat/Catha edulis, cigarettes, and coffee) in an individual's lifetime.

Current substance user. A person who consumed substances (alcohol, khat, cigarettes, and coffee) at least once within the last 30 days.

Substance ever uses. Use of any of the substances (alcohol, khat, cigarettes, and coffee) at least once in an individual's lifetime.

Regular use of computers or television. Reading or watching of computers or television at least once a day for not less than 2 hours [35].

Medication history. Taking of anti-rheumatic, anti-psychiatric & anti-thrombotic drugs in the last one month.

Data collection tools and procedure

Data were collected using a validated interviewer-administered questionnaire comprised of socio-demographic information (age, occupation, educational level, and residence), pregnancy-related characteristics (gestational age, GDM, and parity), and history of medical conditions (anemia, stress, sleep disturbance, and medication history). Visual acuity test (both unaided and with current correction,) was measured by Snellen's illiterate "E" chart in a well-illuminated room, from the women positioned at a 6-meter distance from the chart. Non-cycloplegic refraction was performed using trial lenses, trial frame, and retinoscopy in a semi-dark examination room. Adult sized blood pressure cuff was used to measure systolic and diastolic blood pressure. The data were collected by two BSc Midwives and two Optometrists.

Data quality management

To assure the data quality, high emphasis was given in designing data collection tools. A pre-test was done at Polly Health Center at Gondar town before the actual data collection. The questionnaire was prepared in English and then translated to Amharic (local language) by language expert and retranslated to English by another language expert to check its consistency. Training was also given for the data collectors and supervisor regarding the purpose of the study, ethical issues, and interview and measurement techniques. Strict supervision was undertaken throughout the data collection. The supervisor and principal investigator daily checked the questionnaire for accuracy and completeness.

Data processing and analysis procedure

The collected data were checked for completeness, then entered into EpiData version 3.02 and analyzed using STATA 14. Descriptive measures such as median, frequency, and interquartile range (IQR) were calculated. Both bivariable and multivariable binary logistic regression analyses were performed. Those variables with a p-value of < 0.25 in the bivariable analysis were entered into multivariable logistic regression. We also determined the Adjusted Odds Ratio (AOR) with its 95% confidence interval for those variables included in the multivariable logistic regression model. In the final model, variables with a p-value ≤ 0.05 were declared as significantly associated with RE.

Ethical approval and consent to participate

Ethical clearance with a reference number of 1828/2012 was obtained from the Institutional Review Board (IRB) of the UoG and letter of cooperation was gotten from the UoGCSH. Written informed consent was taken from each study participant. Privacy and confidentiality of information were kept properly and names were not recorded. Those study participants who had severe RE at the time of data collection were referred to the Department of Ophthalmology for further diagnosis and management.

Results

Socio-demographic characteristics of the study participants

A total of 401 pregnant women participated in this study with a response rate of 99.50%. The age range of participants was from 17 to 43 years with a median age of 27 (IQR = 24–31) years. Majority of the respondents were Christianity followers (79.80%) and urban dwellers (83.54%). Regarding educational status, 151 (37.66%) had an education level of college/university ([Table 1](#)).

Clinical, substance use and pregnancy-related characteristics of the study participants

The gestational age of pregnant women ranged from 4 to 40 weeks. Nineteen (4.74%) of the study participants had GDM and about 7% had a history of pregnancy-induced hypertension. Majority of the study participants (63.09%) had a history of contraceptive use prior to their current pregnancy. One-hundred and ninety-six (48.88%) pregnant women had a history of regular use of a computer or watching television for more than 2 hours per day ([Table 2](#)).

Prevalence of refractive error and its associated factors

The overall prevalence of RE among pregnant women in this study was 35.66% (95% CI: 30.95%–40.37%). The most frequent type of RE was myopia (64.34%) followed by hyperopia (31.47%) and antimetropia (4.20%). The prevalence of myopia, hyperopia and antimetropia in both or either eyes from the total study participants was 22.94% (92/401), 11.22% (45/401), and 1.50% (6/401), respectively. Higher prevalence of RE was found among pregnant women with GDM (78.95%), PIH (89.29%), those who regularly used a computer or watched TV (38.78%).

On bivariable logistic regression analysis, RE was associated with the age of study participants, residence, educational status, occupation, income, parity, gestational age, GDM, PIH, family history of spectacle use, regular use of computers or watching TV, medication history, history of contraceptive use, sleep duration, history of lifetime coffee drink, and hematocrit

Table 1. Socio-demographic characteristics of pregnant women attending ANC unit at the University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia, 2020(n = 401).

Variables	Frequency	Percentage (%)
Age in years ^c	Median = 27(IQR = 24–31)	
Religion		
Christian	320	79.80
Muslim	81	20.20
Residence		
Urban	335	83.54
Rural	66	16.46
Educational status		
Cannot read & write	53	13.22
Grade 1–8	88	21.95
Grade 9–12	109	27.18
College/University	151	37.66
Occupation		
Government employee	104	25.94
Private employee	79	19.70
Merchant	40	9.98
Housewife	135	33.67
Others*	43	10.72
Monthly income ^c (ETB)	Median = 3000(IQR = 1200–45000)	

C = continuous variable expressed in median and IQR, ETB = Ethiopian Birr, IQR = interquartile range,

*others = farmers, daily workers and unemployed.

<https://doi.org/10.1371/journal.pone.0246174.t001>

level ($p < 0.25$). However, in the final model, only age, educational level, parity, gestational age and regular use of computers or watching TV were significantly associated with RE ($p \leq 0.05$).

For a unit increase in age of study participants, the odds of developing RE was increased by 1.3 times (AOR = 1.31; 95% CI: 1.16–1.48). RE was increased by 15% (AOR = 1.15, 95% CI: 1.08–1.22) for each week increase of gestational age. The odds of getting RE was 6.2 times (AOR = 6.12, 95% CI: 2.46–15.59) higher among study participants who regularly used computers or watched television than the non-users. Study participants with 9–12 education level had 90% (AOR = 0.10, 95% CI: 0.02–0.54) reduced odds of refractive error than those who cannot read and write. A one-unit increase in the number of parity of study participants led to a 3.20 fold (AOR = 3.17, 95% CI: 1.92–5.25) increase in the odds of developing RE (Table 3).

Discussion

Pregnancy is a physiological condition responsible for a variety of alterations in all body systems including the visual system. These physiological stresses during pregnancy are possibly due to changes in the hormonal, immunological, metabolic, hematologic, and cardiovascular status of the body [44,45]. Even though most of the ocular (refractive) changes are temporary (physiological) that will return to normal after delivery, there are also permanent (pathological) changes persisting after a postpartum period that may require an intervention [46]. To the best of our knowledge, there is no established study on the prevalence of RE among pregnant women in Ethiopia.

In this study, the overall prevalence of RE was 35.66% (95% CI: 30.95%–40.37%) which is lower than studies in India (65%) [11], Nigeria (90%) [10], and South India (77.50%) [12]. This variation might be accounted for the differences in study design used (cross-sectional in our

Table 2. Clinical, substance use and pregnancy-related characteristics of pregnant women attending ANC unit at the University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia, 2020(n = 401).

Variables	Frequency	Percentage (%)
Parity ^c	1(0–2)	
Gestational age ^c (in weeks)	30(24–36)	
Gestational diabetes mellitus		
Yes	19	4.74
No	382	95.26
Pregnancy-induced hypertension		
Yes	28	6.98
No	373	93.02
Family history of spectacle use		
Yes	36	8.98
No	365	91.02
Regular use of computer or television		
Yes	196	48.88
No	205	51.12
Medication history		
Yes	38	9.48
No	363	91.52
Contraceptive use		
Yes	253	63.09
No	148	36.91
Sleep duration		
Short	47	11.72
Optimal	252	62.84
Long	102	25.44
Ever drink alcohol		
Yes	270	67.33
No	131	32.67
Currently drink alcohol		
Yes	176	65.19
No	94	34.81
Ever drink coffee		
Yes	339	84.54
No	62	15.46
Currently drink coffee		
Yes	276	81.42
No	63	18.58
Perceived stress level		
Low	75	18.70
Moderate	294	73.32
High	32	7.98
Hematocrit level ^c (%)	39(37–41)	

C: Continuous variable.

<https://doi.org/10.1371/journal.pone.0246174.t002>

study) and socioeconomic factors. For instance, access to use computers or other digital devices that can precipitate the occurrence of RE in Ethiopia is very low as compared to other studies in developed countries. Therefore, these factors might be responsible for the observed discrepancy.

Table 3. Bivariable and multivariable logistic regression analysis of factors associated with refractive error among study participants attending at ANC unit of University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia, 2020 (n = 401).

Variables	Refractive error		OR (95%CI)	
	Yes N (%)	No N (%)	COR	AOR
Age (years)	31(28–35)	25(23–28)	1.39(1.29–1.48)	1.31(1.16–1.48)*
Residence				
Urban	108(32.24)	227(67.76)	0.42(0.25–0.72)	0.76(0.21–2.77)
Rural	35(53.03)	31(46.97)	1.00	1.00
Educational status				
Can't read & write	32(60.38)	21(39.62)	1.00	1.00
Grade 1–8	31(35.23)	57(64.77)	0.36(0.18–0.72)	0.63(0.17–2.33)
Grade 9–12	11(10.09)	98(89.91)	0.07(0.03–0.17)	0.10(0.02–0.54)*
College/University	69(45.70)	82(54.30)	0.55(0.29–1.04)	0.95(0.19–4.76)
Occupation				
Gov't employee	44(42.31)	60(57.69)	1.62(0.95–2.77)	0.88(0.22–3.51)
Private employee	31(39.24)	48(60.76)	1.43(0.80–2.56)	1.44(0.39–5.28)
Merchant	12(30.00)	28(70.00)	0.95(0.44–2.05)	0.43(0.09–1.95)
House wife	42(31.11)	93(68.89)	1.00	1.00
Others**	14(32.56)	29(67.44)	1.07(0.51–2.23)	1.06(0.20–5.48)
Income (ETB)	4000(2000–5000)	3000(500–4200)	1.00(1.00–1.03)	1.00(0.98–1.00)
Parity	2(2–3)	0(0–1)	4.57(3.39–6.16)	3.17(1.92–5.25)*
Gestational age (week)	34(28–36)	28(20–32)	1.12(1.08–1.16)	1.15(1.08–1.22)*
Gestational DM				
Yes	15(78.95)	4(21.05)	7.44(2.42–22.88)	0.63(0.10–3.92)
No	128(33.51)	254(66.49)	1.00	1.00
PIH				
Yes	25(89.29)	3(10.71)	18.01(5.33–60.83)	2.03(0.39–10.56)
No	118(31.64)	255(68.36)	1.00	1.00
Family history of spectacle use				
Yes	21(58.33)	15(41.67)	2.79(1.39–5.60)	2.30(0.72–7.40)
No	122(33.42)	243(66.58)	1.00	1.00
Regular use of a computer or TV				
Yes	76(38.78)	120(61.22)	1.30(0.87–1.97)	6.19(2.46–15.59)*
No	67(32.68)	138(67.32)	1.00	1.00
Medication history				
Yes	26(68.42)	12(31.58)	4.56(2.22–9.35)	1.91(0.50–7.31)
No	117(32.23)	246(67.77)	1.00	1.00
Contraceptive use				
Yes	131(51.78)	122(48.22)	12.17(6.42–23.07)	2.317(0.817–6.60)
No	12(8.11)	136(91.89)	1.00	1.00
Sleep duration				
Short	26(55.32)	21(44.68)	6.19(2.85–13.45)	1.932(0.49–7.65)
Optimal	100(39.68)	152(60.32)	3.29(1.84–5.87)	0.62(0.25–1.55)
Long	17(16.67)	85(83.33)	1.00	1.00
Ever drink coffee				
Yes	126(37.17)	213(62.83)	1.57(0.86–2.85)	2.16(0.73–6.35)
No	17(27.42)	45(72.58)	1.00	1.00
Hematocrit level ^c (%)	39.4(37.4–41)	38.85(37–40.8)	1.04(0.98–1.11)	0.97(0.87–1.09)

AOR = Adjusted odds ratio, CI = confidence interval, COR = crude odds ratio, ETB = Ethiopian birr, Gov't = government, TV = Television, * = p-value ≤ 0.05,

**others = farmers, daily workers and unemployed.

<https://doi.org/10.1371/journal.pone.0246174.t003>

The prevalence of RE in our study is higher than other studies in Saudi Arabia (25.36%) [8], India (10.50%) [47], Ghana (21.60%) [48], and Rwanda (18.90%) [49]. This discrepancy might be because of the difference in the study population, in which our study population were pregnant women but the general population for other studies. Therefore, the observed higher prevalence of RE in this study might be due to pregnancy-induced endocrine changes. An increased level of estrogen and progesterone during pregnancy is associated with a higher chance of developing refractive errors (either myopia or hyperopia). The reason for this incidence of RE is because of fluid retention in the cornea triggered by estrogen and progesterone. Consequently, fluid retention in the corneal region leads to corneal edema, progressive loss of corneal sensitivity, increased central corneal thickness and curvature, increased thickness of the lens and transient loss of accommodation, and finally end up with RE [39,44,50,51].

The association of maternal age and RE was established in our study. A unit increase in the age of the study participant is associated with a 1.3 times increment of RE. This finding is supported by previous population-based studies in the USA [18], South India [19], India [47], Sri Lanka [52], China [38,53], and Canada [54]. This relationship might be because of an increase in the firmness & hardening of the nucleus of the lens and thickness of cortex of the lens with age secondary to increased insoluble protein molecules and thus, decreases refractive index. With increasing age, the crystalline lens and corneal curvature become steeper and result in RE [18,55,56].

In our study, for each week increase in the gestational age, the odd of having RE was increased by 1.15 times. This finding is comparable with other studies in the USA [57], Iran [23], India [12], and Nigeria [10,24]. The reason for this change might be due to an increased thickness and curvature of the cornea from first to third trimesters of pregnancy following a higher rate of fluid retention triggered by hormones mainly estrogen and progesterone [50]. Aldosterone is also highly secreted during pregnancy and has a synergistic effect with estrogen to cause reabsorption of excess sodium from the renal tubules. This in turn causes the maternal blood volume to rise that leads to accumulation of fluid in the cornea [51]. Other supplemental causes for this incident may also be because of an increased in metabolic rates and high oxygen intake, production of extra red blood cells as a result of increased activities of the bone marrow due to activation of erythropoietin by Placental Lactogen and the rise of cardiac output and heart rate to go with the excess fluid volume [51,58].

A unit increase in the number of births (parity) among pregnant women was associated with increased odds of RE. The finding of our study is in agreement with the finding of a study in Croatia (28) in which, corneal edema, thickness, and curvature were more elevated in multiparous than nulli and primiparous women during pregnancy. The other plausible reason that plays a role for this event is hormonal influences on the successive pregnancies which will upset the physiological refraction system of the eyes.

The likelihood of developing RE among study participants who had a history of regular use of a computer or watching television was higher than those who didn't use. This is fairly consistent with a study in Ethiopia [35], China [59], Iran [1], India [36], and Egypt [60]. This association might be anticipated because staring on the computer, watching television and using other digital devices for a long time (more than 2 hours per day) could induce prolonged accommodation and muscle fatigue (eyestrain) that may result in a transient myopic shift [61,62]. Besides, this positive association might be coined from decreased blink rate and incomplete blinks of eyes while staring on computer or television for a prolonged time. When the eyes are exposed to an incomplete and reduced frequency of blinking during exhaustive use of digital instruments, the tear film doesn't properly run across the whole cornea. This will bring about dry eye symptoms and alter the refractive power of the cornea [62,63].

Our study found that those study participants with an educational level of grade 9–12 were less likely to experience RE than those who couldn't read and write. Contrarily, previous studies

in the United Kingdom [21], Turkey [22], China [64], and Singapore [65] showed a direct association between RE and advanced level of education. The inverse association in our study might be accounted by relatively smaller sample size, less exposure for risk factors (since most of them are housewives), and environmental and socio-economic conditions like low access to use computers or other digital devices as compared to other studies in developed countries.

The present study was cross-sectional and therefore, we are unable to show the cause-effect relationship between independent variables and refractive error. We didn't also apply cycloplegic refraction test to control the accommodation ability of the lens because of its least feasibility in our institution even though its effect is very little. Lastly, we did not collect data regarding sunlight exposure level and time spent in the outdoor which might affect the outcome variable.

Conclusion

The prevalence of RE was high among pregnant women, of which myopia was the most common variety. Advanced age, increased gestational age, increased parity, and regular use of a computer or watching television were factors significantly and positively associated with RE among pregnant women. Apart from providing other maternal health services, a routine screening, and evaluation of pregnant women for RE during ANC visit is also recommended.

Supporting information

S1 File.
(DOCX)

Acknowledgments

We would like to acknowledge the study participants, data collectors and supervisors for their willingness, valuable support and assistance during this work. We are deeply thankful to the staff members of the Department of Optometry at the UoGCSH for their provision of different data collection tools and taking part in the data collection.

Author Contributions

Conceptualization: Mengistie Diress.

Data curation: Mengistie Diress, Yigizie Yeshaw, Minychil Bantihun, Baye Dagne, Adugnaw Ambelu, Mohammed Abdu Seid, Yonas Akalu.

Formal analysis: Mengistie Diress.

Funding acquisition: Mengistie Diress.

Investigation: Mengistie Diress, Yigizie Yeshaw, Minychil Bantihun, Baye Dagne, Adugnaw Ambelu, Mohammed Abdu Seid, Yonas Akalu.

Methodology: Mengistie Diress, Yigizie Yeshaw, Minychil Bantihun, Baye Dagne, Adugnaw Ambelu, Mohammed Abdu Seid, Yonas Akalu.

Project administration: Mengistie Diress.

Resources: Mengistie Diress.

Supervision: Mengistie Diress, Baye Dagne, Adugnaw Ambelu, Yonas Akalu.

Visualization: Yigizie Yeshaw, Minychil Bantihun, Baye Dagne, Adugnaw Ambelu, Mohammed Abdu Seid, Yonas Akalu.

Writing – original draft: Yigizie Yeshaw, Minychil Bantihun, Baye Dagnew, Adugnaw Ambelu, Mohammed Abdu Seid, Yonas Akalu.

Writing – review & editing: Yigizie Yeshaw, Minychil Bantihun, Baye Dagnew, Adugnaw Ambelu, Mohammed Abdu Seid, Yonas Akalu.

References

1. Hashemi H, Fotouhi A, Yekta A, Pakzad R, Ostadimoghaddam H, Khabazkhoob M. Global and regional estimates of prevalence of refractive errors: Systematic review and meta-analysis. *Journal of current ophthalmology*. 2018; 30(1):3–22. <https://doi.org/10.1016/j.joco.2017.08.009> PMID: 29564404
2. Organization WH. Global Initiative for the Elimination of Avoidable Blindness: action plan 2006–2011: Geneva 2017 [cited 2020]. Available from: <https://apps.who.int/iris/handle/10665/43754>.
3. Flaxman SR, Bourne RR, Resnikoff S, Ackland P, Braithwaite T, Cicinelli MV, et al. Global causes of blindness and distance vision impairment 1990–2020: a systematic review and meta-analysis. *The Lancet Global Health*. 2017; 5(12):e1221–e34. [https://doi.org/10.1016/S2214-109X\(17\)30393-5](https://doi.org/10.1016/S2214-109X(17)30393-5) PMID: 29032195
4. Naidoo K, Gichuhi S, Basáñez M-G, Flaxman SR, Jonas JB, Keeffe J, et al. Prevalence and causes of vision loss in sub-Saharan Africa: 1990–2010. *British Journal of Ophthalmology*. 2014; 98(5):612–8. <https://doi.org/10.1136/bjophthalmol-2013-304081> PMID: 24568870
5. Berhane Y, Worku A, Bejiga A, Adamu L, Alemayehu W, Bedri A, et al. Prevalence and causes of blindness and low vision in Ethiopia. *Ethiopian Journal of Health Development*. 2007; 21(3):204–10.
6. Organization WH. Global Initiative for the Elimination of Avoidable Blindness: action plan 2006–2011. 2007.
7. Stevens GA, White RA, Flaxman SR, Price H, Jonas JB, Keeffe J, et al. Global prevalence of vision impairment and blindness: magnitude and temporal trends, 1990–2010. *Ophthalmology*. 2013; 120(12):2377–84. <https://doi.org/10.1016/j.ophtha.2013.05.025> PMID: 23850093
8. Parrey MUR, Elmorsy E. Prevalence and pattern of refractive errors among Saudi adults. *Pakistan journal of medical sciences*. 2019; 35(2):394. <https://doi.org/10.12669/pjms.35.2.648> PMID: 31086521
9. Khalaj M, Mohammadi Zeidi E, Gorbani S, Hashemi F. Assessment of the prevalence of refractive eye error and IOP during pregnancy and after delivery in patients referred to ophthalmology clinic of Boo-Ali Hospital of Qazvin in 1387. *Journal of Mazandaran University of Medical Sciences*. 2010; 19(74):25–31.
10. Nkiru ZN, Obiekwe O, Lilian O, Daniel CN, Uchenna IN, Rich U. Visual acuity and refractive changes among pregnant women in Enugu, Southeast Nigeria. *Journal of family medicine and primary care*. 2018; 7(5):1037–41. https://doi.org/10.4103/jfmpc.jfmpc_335_17 PMID: 30598953
11. Balasubramanian K MS, Nagarajan G. A. Prospective Study of Changes in the Refractive System of Eye during Pregnancy. *INTERNATIONAL JOURNAL OF SCIENTIFIC STUDY*. 2017; 5(4):89–92.
12. Usha C PB, Kalaiselvi B. A study on changes in the central corneal thickness among pregnant mothers in south India. *Indian Journal of Clinical and Experimental Ophthalmology*. 2019; 5(2):219–21.
13. Resnikoff S, Pascolini D, Mariotti SP, Pokharel GP. Global magnitude of visual impairment caused by uncorrected refractive errors in 2004. *Bulletin of the World Health Organization*. 2008; 86(1):63–70. <https://doi.org/10.2471/blt.07.041210> PMID: 18235892
14. Ofir D, Kessous R, Belfer N, Lifshitz T, Sheiner E. The influence of visual impairment on pregnancy outcomes. *Archives of gynecology and obstetrics*. 2015; 291(3):519–23. <https://doi.org/10.1007/s00404-014-3412-4> PMID: 25141991
15. Trilaksana Nugroho HPJ, Dea Amarilisa Adespin, Bambang Hariyana, Ari Budi Himawan. ASSOCIATION BETWEEN REFRACTIVE ERROR AND DECREASING VISUS IN PREGNANT WOMEN. *International conference on Public health 2018*:288.
16. George Iancu VC, Iancu Raluca, Corbu Catalina. Particularities of myopia in pregnancy. *Gineco+EU*. 2013; 9(34).
17. E SK. *The Eye: The Physiology of Human Perception*. The Rosen Publishing Group, Inc 2010.
18. Richdale K, Bullimore MA, Sinnott LT, Zadnik K. The Effect of Age, Accommodation, and Refractive Error on the Adult Human Eye. *Optometry and vision science: official publication of the American Academy of Optometry*. 2016; 93(1):3–11.
19. Krishnaiah S, Srinivas M, Khanna RC, Rao GN. Prevalence and risk factors for refractive errors in the South Indian adult population: The Andhra Pradesh Eye disease study. *Clinical ophthalmology (Auckland, NZ)*. 2009; 3:17–27. PMID: 19668540

20. Rai BB, Ashby RS, French AN, Maddess T. Rural-urban differences in myopia prevalence among myopes presenting to Bhutanese retinal clinical services: a 3-year national study. *Graefes's Archive for Clinical and Experimental Ophthalmology*. 2020;1–9. <https://doi.org/10.1007/s00417-020-04891-6> PMID: 32803328
21. Mountjoy E, Davies NM, Plotnikov D, Smith GD, Rodriguez S, Williams CE, et al. Education and myopia: assessing the direction of causality by mendelian randomisation. *bmj*. 2018; 361:k2022. <https://doi.org/10.1136/bmj.k2022> PMID: 29875094
22. Ilhan C. Investigation of the Refractive Errors in Terms of Education Level in Central Anatolian Population. *EC Ophthalmology*. 2018; 9:308–11.
23. Mehdizadehkashi K, Chaichian S. Corrigendum to "Visual Acuity Changes during Pregnancy and Postpartum: A Cross-Sectional Study in Iran". 2018; 2018:5280697.
24. Ebeigbe JA, Ebeigbe PN, Ighoroje A. Ocular changes in pregnant Nigerian women. *Nigerian journal of clinical practice*. 2012; 15(3):298–301. <https://doi.org/10.4103/1119-3077.100624> PMID: 22960964
25. Sabel BA, Wang J, Cárdenas-Morales L, Faiq M, Heim C. Mental stress as consequence and cause of vision loss: the dawn of psychosomatic ophthalmology for preventive and personalized medicine. *EPMA journal*. 2018; 9(2):133–60. <https://doi.org/10.1007/s13167-018-0136-8> PMID: 29896314
26. Jee D, Morgan IG, Kim EC. Inverse relationship between sleep duration and myopia. *Acta ophthalmologica*. 2016; 94(3):e204–e10. <https://doi.org/10.1111/aos.12776> PMID: 26031352
27. Stambolian D. Genetic susceptibility and mechanisms for refractive error. *Clinical genetics*. 2013; 84(2):102–8. <https://doi.org/10.1111/cge.12180> PMID: 23647423
28. Shaaban YM, Badran TAF. The effect of oral contraceptive pills on the macula, the retinal nerve fiber layer, the ganglion cell layer and the choroidal thickness. *BMC ophthalmology*. 2019; 19(1):250. <https://doi.org/10.1186/s12886-019-1263-2> PMID: 31823761
29. Venkataraman A, Panda BB, Anupam Dey M. Nutritional anemia as a cause of vision loss in developing countries: A case report. *Kerala journal of ophthalmology*. 2013; 25(4):383–4.
30. Yarbağ A, Yazar H, Akdoğan M, Pekgör A, Kaleli S. Refractive errors in patients with newly diagnosed diabetes mellitus. *Pakistan journal of medical sciences*. 2015; 31(6):1481. <https://doi.org/10.12669/pjms.316.8204> PMID: 26870120
31. Toygar O, Sizmaz S, Pelit A, Toygar B, KIZILOĞLU ÖY, Akova Y. Central corneal thickness in type II diabetes mellitus: is it related to the severity of diabetic retinopathy? *Turkish journal of medical sciences*. 2015; 45(3):651–4. <https://doi.org/10.3906/sag-1404-153> PMID: 26281334
32. Rezaei S, LoBue S, LoBue T, Giovane R, Carney M, Henderson C. Ophthalmic complications and ocular changes in pregnancy-A Review. *Obstet Gynecol Int J*. 2016; 4(1):00093.
33. Roos NM, Wiegman MJ, Jansonius NM, Zeeman GG. Visual disturbances in (pre) eclampsia. *Obstetrical & gynecological survey*. 2012; 67(4):242–50. <https://doi.org/10.1097/OGX.0b013e318250a457> PMID: 22495060
34. Kaimbo D. Transient drug-induced myopia. *SAJ Case Rep*. 2014; 1(1):105.
35. Sewunet SA AK, Gedefew M. Uncorrected refractive error and associated factors among primary school children in Debre Markos District, Northwest Ethiopia. *BMC ophthalmology*. 2014; 14(1):95. <https://doi.org/10.1186/1471-2415-14-95> PMID: 25070579
36. Kumar KS AB. Prevalence of refractive error among school-going children of Imphal Manipur. *Int J Med Sci Public Health*. 2016; 5:1364–8.
37. Andualem HB, Assefa NL, Weldemichael DZ, Tefera TK. Prevalence and associated factors of presbyopia among school teachers in Gondar city, Northwest Ethiopia, 2016. *Clinical optometry*. 2017; 9:85–90. <https://doi.org/10.2147/OPTO.S129326> PMID: 30214364
38. Wang M, Cui J, Shan G, Peng X, Pan L, Yan Z, et al. Prevalence and risk factors of refractive error: a cross-sectional Study in Han and Yi adults in Yunnan, China. *BMC ophthalmology*. 2019; 19(1):33. <https://doi.org/10.1186/s12886-019-1042-0> PMID: 30683073
39. Sharma S, Rekha W, Sharma T, Downey G. Refractive issues in pregnancy. *The Australian & New Zealand journal of obstetrics & gynaecology*. 2006; 46(3):186–8. <https://doi.org/10.1111/j.1479-828X.2006.00569.x> PMID: 16704469
40. S. M. Antimetropia in a 10-year-old boy with unilateral tilted disc syndrome. *Clinical Optometry* 2014; 6(43).
41. Cappellini MD, Motta I. Anemia in Clinical Practice-Definition and Classification: Does Hemoglobin Change With Aging? *Seminars in hematology*. 2015; 52(4):261–9. <https://doi.org/10.1053/j.seminhematol.2015.07.006> PMID: 26404438
42. Peltzer K, Phaswana-Mafuya N. Association between Visual Impairment and Low Vision and Sleep Duration and Quality among Older Adults in South Africa. *International journal of environmental research and public health*. 2017;14(7). <https://doi.org/10.3390/ijerph14070811> PMID: 28753944

43. Jee D, Morgan IG, Kim EC. Inverse relationship between sleep duration and myopia. *Acta ophthalmologica*. 2016; 94(3):e204–10. <https://doi.org/10.1111/aos.12776> PMID: 26031352
44. Kalogeropoulos D, Sung VC. The physiologic and pathologic effects of pregnancy on the human visual system. 2019; 39(8):1037–48.
45. Gotovac M KS, Lukenda A. Eye and Pregnancy. *Collegium antropologicum*. 2013; 37(1):189–93.
46. Samra KA. The eye and visual system in pregnancy, what to expect? An in-depth review. *Oman journal of ophthalmology*. 2013; 6(2):87–91. <https://doi.org/10.4103/0974-620X.116626> PMID: 24082665
47. Karavadi Sri Sai Vidusha DMN. Prevalence of refractive errors among school children in the rural field practice area of a tertiary care hospital, Bengaluru. *International Journal of Community Medicine and Public Health*. 2018; 5(4):1471–6.
48. DO A. Prevalence and Causes of Visual Impairment among Patients in Juaben Hospital Eye Clinic, Ghana. *M J Opht*. 2017; 2(2):017.
49. Semanyenzi SE KJ, Nzayirambaho M. Prevalence and pattern of refractive errors in high schools of Nyarugenge district. *Rwanda Medical Journal*. 2015; 72(3):8–13.
50. Agrawal N, Agarwal LT, Lavaju P, Chaudhary SK. Physiological Ocular Changes in Various Trimesters of Pregnancy. *Nepalese journal of ophthalmology: a biannual peer-reviewed academic journal of the Nepal Ophthalmic Society: NEPJOPH*. 2018; 10(19):16–22. <https://doi.org/10.3126/nepjoph.v10i1.21685> PMID: 31056572
51. Sanghavi M, Rutherford JD. Cardiovascular physiology of pregnancy. *Circulation*. 2014; 130(12):1003–8. <https://doi.org/10.1161/CIRCULATIONAHA.114.009029> PMID: 25223771
52. Gilbert C MG, Schmidt E, Edussuriya K, Kumara RP, Wimalaratne SA, Athapattu AH, et al. Prevalence and types of refractive errors, and spectacle coverage in Sri Lankan adults: The Sri Lanka National survey of blindness and visual impairment. *Ceylon Medical Journal* 2018 63(5):33.
53. Xu L, Wang Y, Li Y, Wang Y, Cui T, Li J, et al. Causes of blindness and visual impairment in urban and rural areas in Beijing: the Beijing Eye Study. *Ophthalmology*. 2006; 113(7):1134.e1–11. <https://doi.org/10.1016/j.ophtha.2006.01.035> PMID: 16647133
54. Irving EL, Machan CM, Lam S, Hrynchak PK, Lillakas L. Refractive error magnitude and variability: Relation to age. *Journal of optometry*. 2019; 12(1):55–63. <https://doi.org/10.1016/j.optom.2018.02.002> PMID: 29567041
55. Skalkicky SE. *Ocular and Visual Physiology*. Springer Science+Business Media Singapore 2016.
56. Richdale K, Sinnott LT, Bullimore MA, Wassenaar PA, Schmalbrock P, Kao CY, et al. Quantification of age-related and per diopter accommodative changes of the lens and ciliary muscle in the emmetropic human eye. *Investigative ophthalmology & visual science*. 2013; 54(2):1095–105.
57. Wu F, Schallhorn JM, Lowry EA. Refractive status during pregnancy in the United States: results from NHANES 2005–2008. *Graefe's archive for clinical and experimental ophthalmology = Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie*. 2020; 258(3):663–7. <https://doi.org/10.1007/s00417-019-04552-3> PMID: 31823061
58. Lee LK. Physiological adaptations of pregnancy affecting the nervous system. *Seminars in neurology*. 2007; 27(5):405–10. <https://doi.org/10.1055/s-2007-991124> PMID: 17940919
59. Guan H, Yu NN. Impact of various types of near work and time spent outdoors at different times of day on visual acuity and refractive error among Chinese school-going children. 2019; 14(4):e0215827.
60. Saad A, El-Bayoumy BM. Environmental risk factors for refractive error among Egyptian schoolchildren. *Eastern Mediterranean health journal = La revue de sante de la Mediterranee orientale = al-Majallah al-sihhiyah li-sharq al-mutawassit*. 2007; 13(4):819–28. PMID: 17955764
61. Kim SH, Suh YW, Choi YM, Han JY, Nam GT, You EJ, et al. Effect of watching 3-dimensional television on refractive error in children. *Korean journal of ophthalmology: KJO*. 2015; 29(1):53–7. <https://doi.org/10.3341/kjo.2015.29.1.53> PMID: 25646061
62. Sameena Kokab MIK. COMPUTER VISION SYNDROME: A SHORT REVIEW. *Journal of Evolution of Medical and Dental Sciences*. 2012; 1(6):1223.
63. Alemayehu AM AM. Pathophysiologic Mechanisms of Computer Vision Syndrome and its Prevention. *World J Ophthalmol Vis Res*. 2019; 2(5):1–7.
64. Wang M, Cui J, Shan G, Peng X, Pan L, Yan Z, et al. Prevalence and risk factors of refractive error: a cross-sectional Study in Han and Yi adults in Yunnan, China. *BMC ophthalmology*. 2019; 19(1):33. <https://doi.org/10.1186/s12886-019-1042-0> PMID: 30683073
65. Tan C, Chan Y, Wong TY, Gazzard G, Niti M, Ng TP, et al. Prevalence and risk factors for refractive errors and ocular biometry parameters in an elderly Asian population: the Singapore Longitudinal Aging Study (SLAS). *Eye*. 2011; 25(10):1294–301. <https://doi.org/10.1038/eye.2011.144> PMID: 21720418